

Cheat Sheet Template

Physical Layer

$$\frac{dT}{dt} = -k(T - T_o)$$

T_o = outside temperature



A colourful spiral

Inner Product Spaces

- $\langle v, v \rangle \geq 0$ Furthermore, $\langle v, v \rangle = 0 \leftrightarrow v = 0$
 - $\langle v, u \rangle = \langle u, v \rangle$
 - $\langle ku, v \rangle = k\langle u, v \rangle$
 - $\langle u + v, w \rangle = \langle u, w \rangle + \langle v, w \rangle$
- $\|v\| = \sqrt{\langle v, v \rangle}$
 $\cos^{-1}\left(\frac{\langle v, u \rangle}{\|v\|\|u\|}\right)$

Gram-Schmidt

$$v_1 = x_1$$

$$v_2 = x_2 - \frac{\langle x_2, v_1 \rangle}{\|v_1\|^2} v_1$$

$$\vdots$$

$$v_n = x_n - \sum_{k=1}^{n-1} \frac{\langle x_n, v_k \rangle}{\|v_k\|^2} v_k$$

Variation of Parameters

$$F(x) = y'' + y'$$

$y_h = b_1 y_1(x) + b_2 y_2(x)$, y_1, y_2 are L.R.

$$y_p = u_1(x)y_1(x) + u_2(x)y_2(x)$$

$$u_1 = \int^t \frac{-y_2 F(t) dt}{w[y_1, y_2](t)}$$

$$u_2 = \int^t \frac{y_1 F(t) dt}{w[y_1, y_2](t)}$$

$$y = y_h + y_p$$

Systems

$$\vec{x}' = A\vec{x}$$

A is diagonalizable $\vec{x}(t) = a_1 e^{\lambda_1 t} \vec{v}_1 + \dots + a_n e^{\lambda_n t} \vec{v}_n$

A is not diagonalizable $\vec{x}(t) = a_1 e^{\lambda_1 t} \vec{v}_1 + a_2 e^{\lambda t} (\vec{w} + t\vec{v})$
 where $(A - \lambda I)\vec{w} = \vec{v}$
 \vec{v} is an Eigenvector w/ value λ
 i.e. \vec{w} is a generalized Eigenvector

$$\vec{x}' = A\vec{x} + \vec{B}$$

Solve y_h
 $\vec{x}_1 = e^{\lambda_1 t} \vec{v}_1, \vec{x}_2 = e^{\lambda_2 t} \vec{v}_2$
 $\vec{X} = [\vec{x}_1, \vec{x}_2]$
 $\vec{X}\vec{u}' = \vec{B}$
 $y_p = \vec{X}\vec{u}$
 $y = y_h + y_p$

ODEs

1st Order Linear	Use integrating factor, $I = e^{\int P(x)dx}$
Separable:	$\int P(y)dy/dx = \int Q(x)$
HomogEnEous:	$dy/dx = f(x, y) = f(xt, yt)$ sub $y = xV$ solve, then sub $V = y/x$
Exact:	If $M(x, y) + N(x, y)dy/dx = 0$ and $M_y = N_x$ i.e. $\langle M, N \rangle = \nabla F$ then $\int_x M + \int_y N = F$
Order Reduction	Let $v = dy/dx$ then check other types If purely a function of y , $\frac{dv}{dx} = v \frac{dv}{dy}$
Variation of Parameters:	When $y'' + a_1 y' + a_2 y = F(x)$ F contains $\ln x$, $\sec x$, $\tan x$, \div
Bernoulli	$y' + P(x)y = Q(x)y^n$ $\div y^n$ $y^{-n} y' + P(x)y^{1-n} = Q(x)$ Let $U(x) = y^{1-n}(x)$ $\frac{dU}{dx} = (1-n)y^{-n} \frac{dy}{dx}$ $\frac{1}{1-n} \frac{dU}{dx} + P(x)U(x) = Q(x)$ solve as a 1st order
Cauchy-Euler	$x^n y^n + a_1 x^{n-1} y^{n-1} + \dots + a_{n-1} y^{n-2} + a_n y = 0$ guess $y = x^r$
3 Cases:	
1) Distinct real roots	$y = ax^{r1} + bx^{r2}$
2) Repeated real roots	$y = Ax^r + y_2$ Guess $y_2 = x^r u(x)$ Solve for $u(x)$ and choose one ($A = 1, C = 0$)
3) Distinct complex roots	$y = B_1 x^a \cos(b \ln x) + B_2 x^a \sin(b \ln x)$

Matrix Exponentiation

$$A^n = SD^n S^{-1}$$

D is the diagonalization of A

Complex Numbers

Systems of equations If $\vec{w}_1 = u(\vec{t}) + iv(\vec{t})$ is a solution,
 $\vec{x}_1 = u(\vec{t}), \vec{x}_2 = v(\vec{t})$ are solutions
 i.e. $\vec{x}_h = c_1 \vec{x}_1 + c_2 \vec{x}_2$

Euler's Identity $e^{ix} = \cos x + i \sin x$

Laplace Transforms

$$L[f](s) = \int_0^\infty e^{-sx} f(x) dx$$

$f(t) = t^n, n \geq 0$	$F(s) = \frac{n!}{s^{n+1}}, s > 0$
$f(t) = e^{at}, a \text{ constant}$	$F(s) = \frac{1}{s-a}, s > a$
$f(t) = \sin bt, b \text{ constant}$	$F(s) = \frac{b}{s^2 + b^2}, s > 0$
$f(t) = \cos bt, b \text{ constant}$	$F(s) = \frac{s}{s^2 + b^2}, s > 0$
$f(t) = t^{-1/2}$	$F(s) = \frac{\pi}{s^{1/2}}, s > 0$
$f(t) = \delta(t - a)$	$F(s) = e^{-as}$
f'	$L[f'] = sL[f] - f(0)$
f''	$L[f''] = s^2 L[f] - sf(0) - f'(0)$
$L[e^{at} f(t)]$	$L[f](s - a)$
$L[u_a(t) f(t - a)]$	$L[f]e^{-as}$

Gaussian Integral

$$\int_{-\infty}^{+\infty} e^{-1/2(\vec{x}^T A \vec{x})} = \frac{\sqrt{2\pi}^n}{\sqrt{\det A}}$$

Vector Spaces

- $v_1, v_2 \in V$
- $v_1 + v_2 \in V$
 - $k \in \mathbb{F}, kv_1 \in V$
 - $v_1 + v_2 = v_2 + v_1$
 - $(v_1 + v_2) + v_3 = v_1 + (v_2 + v_3)$
 - $\forall v \in V, 0 \in V \mid 0 + v_1 = v_1 + 0 = v_1$
 - $\forall v \in V, \exists -v \in V \mid v + (-v) = (-v) + v = 0$
 - $\forall v \in V, 1 \in \mathbb{F} \mid 1 * v = v$
 - $\forall v \in V, k, l \in \mathbb{F}, (kl)v = k(lv)$
 - $\forall k \in \mathbb{F}, k(v_1 + v_2) = kv_1 + kv_2$
 - $\forall v \in V, k, l \in \mathbb{F}, (k+l)v = kv + lv$